

Short communication

### THE PPAR $\alpha$ PATHWAY IN V $\gamma$ 9V $\delta$ 2 T CELL ANERGY

MARY POUPOT<sup>1,2,3,\*</sup>, FRÉDÉRIC BOISSARD<sup>1,2,3</sup>, DELPHINE BETOUS<sup>1,2,3</sup>,  
 LAURE BARDOUILLET<sup>4</sup>, SÉVERINE FRUCHON<sup>5</sup>, FATIMA L'FAQIHI-  
 OLIVE<sup>5</sup>, FRÉDÉRIC PONT<sup>1</sup>, MOURAD MEKAOUCHÉ<sup>4</sup>, SOPHIE  
 INGOURE<sup>6</sup>, HÉLÈNE SICARD<sup>6</sup>, GUY DUBREUILH<sup>4</sup>  
 and JEAN-JACQUES FOURNIÉ<sup>1,2,3</sup>

<sup>1</sup>INSERM UMR1037-Cancer Research Center of Toulouse, <sup>2</sup>ERL 5294 CNRS,  
 BP3028, CHU Purpan, F-31300 Toulouse, France, <sup>3</sup>Université Toulouse III  
 Paul-Sabatier, 31300 Toulouse, France, <sup>4</sup>Station de Primatologie, UPS 846,  
 Centre National de la Recherche Scientifique, F-13790 Rousset sur Arc, France,  
<sup>5</sup>INSERM U1043-CPTP Toulouse, F-31300 France, <sup>6</sup>Innate Pharma,  
 F-13276 Marseille, France

**Abstract:** Phosphoantigens (PAgs) activate V $\gamma$ 9V $\delta$ 2 T lymphocytes, inducing their potent and rapid response *in vitro* and *in vivo*. However, humans and non-human primates that receive repeated injections of PAgs progressively lose their V $\gamma$ 9V $\delta$ 2 T cell response to them. To elucidate the molecular mechanisms of this *in vivo* desensitization, we analyzed the transcriptome of circulating V $\gamma$ 9V $\delta$ 2 T cells from macaques injected with PAg. We showed that three PAg injections induced the activation of the PPAR $\alpha$  pathway in V $\gamma$ 9V $\delta$ 2 T cells. Thus, we analyzed the *in vitro* response of V $\gamma$ 9V $\delta$ 2 T cells stimulated with a PPAR $\alpha$  agonist. We demonstrated that *in vitro* PPAR $\alpha$  pathway activation led to the inhibition of the BrHPP-induced activation and proliferation of human V $\gamma$ 9V $\delta$ 2 T cells. Since the PPAR $\alpha$  pathway is involved in the antigen-selective desensitization of human V $\gamma$ 9V $\delta$ 2 T cells, the use of PPAR $\alpha$  inhibitors could enhance cancer immunotherapy based on V $\gamma$ 9V $\delta$ 2 T cells.

**Keywords:** Activation, Gamma-delta T-lymphocyte, Immunotherapy, Phosphoantigen, TCR, PPAR $\alpha$

---

\* Author for correspondence. Email: [mary.poupot@inserm.fr](mailto:mary.poupot@inserm.fr), phone: +335 8274 1662

Abbreviations used: CFSE – carboxyfluorescein succidimidyl ester; GSEA – *Gene Set Enrichment Analysis*; IL-2 – interleukin 2; PAg – phosphoantigen; PPAR – peroxisome proliferator-activated receptors

## INTRODUCTION

The  $\gamma\delta$  T lymphocytes present hallmarks of both innate and adaptive immunity. In human and non-human primates, the peripheral repertoire of  $\gamma\delta$  TCR is mostly restricted to TCRV $\gamma$ 9V $\delta$ 2, presumably as a result of repeated exposure to few antigens called phosphoantigens (PAGs). Upon activation with PAGs, these cells release pro-inflammatory chemokines and cytokines and mediate cell cytotoxicity, prior to undergoing an intense proliferation leading to clonal expansion. In parallel, V $\gamma$ 9V $\delta$ 2 T cells mature from naive through central memory to Th1 effector memory and cytotoxic TEMRa cells. TEMRa cells mediate potent natural killer-like cytotoxic responses against tumor cell targets [1, 2].

These features make V $\gamma$ 9V $\delta$ 2 T lymphocytes versatile and attractive candidates for new cancer immunotherapy [3–6]. Synthetic analogues of natural PAGs, such as bromohydrin pyrophosphate (BrHPP), have been produced and used in preclinical and clinical studies. They show the potent and rapid response of the V $\gamma$ 9V $\delta$ 2 T lymphocytes in vitro and in vivo. Their response indeed comprises an IL-2-dependent clonal expansion, but surprisingly does not show memory to prior exposure to PAG stimulus.

In a preclinical study of PAG safety in the cynomolgus, the first infusions of BrHPP with low dosages of IL-2 induced a transient amplification and cytokine response of circulating V $\gamma$ 9V $\delta$ 2 T cells, while further infusions of BrHPP and IL-2 led to their progressive exhaustion [7]. In a TB vaccine setting, eight out of eight macaque monkeys that were injected three times with BrHPP in a subunit vaccine responded to the first injection but did not respond to subsequent boosters [8]. Several other unpublished studies from our and other laboratories have confirmed that non-human primates repeatedly injected with PAGs progressively lose their V $\gamma$ 9V $\delta$ 2 T cell response to such synthetic stimuli. In humans, repeated injections of clinical-grade BrHPP (IPH1101, Innate Pharma) also progressively reduce the response of circulating V $\gamma$ 9V $\delta$ 2 T cell [9].

PPARs are nuclear receptors and transcription factors for genes regulating cellular differentiation, development, metabolism and tumorigenesis. PPAR $\alpha$  is mostly expressed in tissues with high rates of mitochondrial fatty acid oxidation. Upon binding of natural unsaturated fatty acids, leukotriene B4 or synthetic fibrates, PPAR $\alpha$  promotes lipid mobilization and catabolism [10]. By contrast, in immune cells PPAR $\alpha$  primarily inhibits inflammatory pathways by repressing c-jun and NF- $\kappa$ B [11, 12]. PPAR $\alpha$  also inhibits IFN $\gamma$  responses of human  $\alpha\beta$  T cells [13]. Whether PPAR $\alpha$  also regulates human  $\gamma\delta$  T cell functions remains unclear.

PAG-selective desensitization has been identified as a pitfall in the development of V $\gamma$ 9V $\delta$ 2 T cell-based cancer immunotherapy [14, 15]. As the first step to elucidate the molecular mechanisms underlying this in vivo desensitization, we analyzed the transcriptome of circulating V $\gamma$ 9V $\delta$ 2 T cells from macaques injected with BrHPP and control subjects that did not receive the injection. The comparison of the transcriptome before injection versus that after three injections

outlined a gene expression pattern (GEP) reflecting activation of the PPAR $\alpha$  pathway. As these results suggested that in vivo PAg desensitization of V $\gamma$ 9V $\delta$ 2 T lymphocytes is associated with activation of PPAR $\alpha$ , we tested whether activation of the PPAR $\alpha$  pathway in human V $\gamma$ 9V $\delta$ 2 T cells affected their response to PAg. We found that PPAR $\alpha$  agonist-treated V $\gamma$ 9V $\delta$ 2 T cells are unresponsive to BrHPP, confirming that activation of PPAR $\alpha$  could contribute to  $\gamma\delta$  cell desensitization.

## MATERIALS AND METHODS

### Primate cell samples

Animal care and handling was carried out at the Experimentation Animal-Station de Primatologie-UPS 846 CNRS (Rousset sur Arc, France) in accordance with standard guidelines, following a protocol submitted to and approved by the local ethical committee. Four age- and sex-matched cynomolgus macaques (2.5–4.4 kg) received three successive injections of BrHPP (doses of 50 mg/kg in 30 min intravenous infusions on days 0, 26 and 55) over 2 months, plus subcutaneous injection of rhIL-2 ( $10^6$  IU, Proleukin Chiron) daily for 5 days after each BrHPP injection. Then, 4 ml of blood were collected prior to the first injection and seven days after each injection of BrHPP, i.e. on day 0 (I1), day 28 (I2) and day 55 (I3). TCRV $\gamma$ 9<sup>+</sup> cells were sorted (> 98%) as described previously [7, 8].

### Microarray procedures

Total RNA from the specified macaque TCRV $\gamma$ 9<sup>+</sup> cell samples was isolated using TRIzol Reagent (Invitrogen Life Technologies). The quality of each RNA was assessed with Agilent 2100 Bioanalyser (Agilent Technologies, Palo Alto, CA, USA). Microarray analyses were performed using 1–3  $\mu$ g total RNA, GeneChip Rhesus Macaque Genome Array and GeneChip Operating Software (Affymetrix). The raw transcriptome data were normalized in batch using the RMA software to yield 27755 annotated probe sets for further comparative analysis. The genes differentially expressed were defined using two-way Student's tests ( $p < 0.01$ ), then analyzed for enrichment in functionally related genes among lists downloaded from the GSEA C1–C5 gene sets collection. Gene-selective enrichment analysis was performed as described previously [16] using the Autocompare freeware (<https://sites.google.com/site/fredsoftwares/products/data-mining>).

### V $\gamma$ 9V $\delta$ 2 T lymphocytes early activation and IFN $\gamma$ production

PBMCs isolated from healthy donors (Etablissement Français du Sang, Toulouse, France) after Ficoll-Hypaque density centrifugation, were cultured at 2.5 M/ml in complete RPMI 1640 medium (Invitrogen), e.g. with L-glutamine, streptomycin, penicillin and sodium-pyruvate (Cambrex Biosciences) supplemented with 10% Fetal Clone I (HyClone/Thermo Fisher Scientific). LY-171883 (0–100  $\mu$ M; Sigma-Aldrich) and 100 nM BrHPP (Innate Pharma) were added 24 h later. CD69 (Beckman Coulter) expression by V $\gamma$ 9V $\delta$ 2 T lymphocytes was measured 48 h later via flow cytometry.

IFN $\gamma$  expression was measured via flow cytometry with a staining in PBS 1% saponin with anti-TCRV $\gamma$ 9 (Biolegend, San Diego, CA) and anti-IFN $\gamma$  (BD Biosciences, San Jose, CA) after 6 h of treatment with 10  $\mu$ g/ml brefeldin A (Sigma-Aldrich) and fixation with PBS 2% PFA.

#### **V $\gamma$ 9V $\delta$ 2 T lymphocytes proliferation**

PBMCs were labeled with 0.5  $\mu$ M CFSE (Invitrogen Life Technologies) for 8 min at 37°C, washed and cultured with or without LY-171883 (0–100  $\mu$ M) and 10 or 100 U/ml rhIL-2 (Sanofi-Aventis). Cells were stimulated 24 h later with 100 nM BrHPP. After six days of culture, cells were stained for TCR V $\gamma$ 9 and analyzed for CFSE dilution via flow cytometry.

### **RESULTS AND DISCUSSION**

Repeated stimulation with PAg selectively desensitizes the V $\gamma$ 9V $\delta$ 2 T cell lineage in humans and non-human primates [7, 8, 17–19]. To investigate this desensitization, we first analyzed the GEP of purified macaque V $\gamma$ 9V $\delta$ 2 T cells from subjects injected with BrHPP. Four cynomolgus macaques received three intramuscular injections of BrHPP on day 0 (I1), day 28 (I2) and day 55 (I3), a regimen which abrogates their V $\gamma$ 9V $\delta$ 2 T cell response to BrHPP [7, 8]. Each blood sample was collected prior to the first injection and seven days after each BrHPP injection. Fresh V $\gamma$ 9V $\delta$ 2 T cells were then purified: their mRNA fraction was isolated, amplified and hybridized to Affymetrix Rhesus microarrays.

The mean of V $\gamma$ 9V $\delta$ 2 T cell transcriptomes from the four macaques at the same time points are referred below to as resting, I1, I2 and I3. The I1, I2 and I3 transcriptomes were respectively compared to the resting one (Fig. 1A and B).

The I1 transcriptome encompassed 1013 genes differentially expressed relative to the resting cells (695 genes upregulated and 317 genes downregulated,  $p < 0.05$ ). Of these 1013 genes, only 483 were fully annotated in macaques. Furthermore, among these 483 macaque I1  $\gamma\delta$  response genes, 146 (30%) are orthologs of human V $\gamma$ 9V $\delta$ 2 T response genes modulated one week after a single in vitro exposure to BrHPP [8].

The I2 transcriptome encompassed 959 genes differentially expressed relative to the resting cells (497 genes upregulated and 462 genes downregulated,  $p < 0.05$ ). Only 198 genes (~20%) were common to the I1 and I2 transcriptomes.

The I3 transcriptome showed a still-distinct pattern of expression, since it comprised only 522 genes significantly modulated (285 upregulated and 237 downregulated,  $p < 0.05$ ), ~80% of which were not in the I1 or I2 transcriptomes. Actually, only 37 genes (3% of the initial transcriptional response) were shared by the I1, I2 and I3 transcriptomes, confirming a global change in the V $\gamma$ 9V $\delta$ 2 T cell gene expression response according to the number of injection cycles (Fig. 1B). Of the modulated 522 I3 genes, 130 were fully annotated in the macaque genome.




Fig. 1. Gene signature of macaque V $\gamma$ 9V $\delta$ 2 T cells repeatedly injected with phosphoantigen unveils induction of the PPAR $\alpha$  pathway. A – The mean of the gene expression level and fold change expression for the I1, I2 and I3 of BrHPP injections respectively compared to the resting one (four macaques). B – Shared genes differentially expressed compared to resting cells for the three injections I1, I2 and I3.

When analyzed with Autocompare software for significant enrichment of gene signatures, the I3 genes NPDC1, GPNMB, GNRH1, SFN (*p14-3-3 $\sigma$*  repressor), EML4, RB1, CDKN2C (*p18*), and PRKCA corresponded to negative regulation of the cell cycle pathway ( $p = 3 \times 10^{-4}$ ). No selective enrichment in T cell anergy or lymphocyte exhaustion gene signatures was found for the I3 genes. Nevertheless, the only other significant gene signature of the I3 V $\gamma$ 9V $\delta$ 2 T cell transcriptome was the PPAR $\alpha$  pathway ( $p = 5 \times 10^{-4}$ , Biocarta) based on upregulation of the EP300, NCOR2, MED1, PRKC1, HSD17B4, PRKAR2B and RB1 genes, while RARA and PRKCA were downregulated. The above results showed a global decrease in transcriptional V $\gamma$ 9V $\delta$ 2 T cell responses to successive in vivo stimulations by PAg and an activation of the PPAR $\alpha$  pathway in macaque V $\gamma$ 9V $\delta$ 2 T cells after a I3 exposure to stimulating PAg.




Fig. 2. PPAR $\alpha$  agonist inhibits CD69 induction, IFN $\gamma$  expression and proliferation of V $\gamma$ 9V $\delta$ 2 T cells. A – Expression of CD69 by V $\gamma$ 9V $\delta$ 2 T cells gated after 2 days of culture with or without BrHPP and LY-171883 (1, 10 or 100  $\mu$ M). B – Mean of percentage of CD69 positive cells from 5 independent experiments. C – Expression of intracellular IFN $\gamma$  by V $\gamma$ 9V $\delta$ 2 T cells cultured with or without LY-171883 (1, 10 or 100  $\mu$ M), mean from 5 independent experiments. D – Expression of intracellular IFN $\gamma$  by V $\gamma$ 9V $\delta$ 2 T cells, one of the 5 experiments. E – Proliferation of V $\gamma$ 9V $\delta$ 2 T cells visualized by CFSE dilution in V $\gamma$ 9V $\delta$ 2 T cells after 6 days of culture with or without LY-171883 (1, 10 or 100  $\mu$ M) and with 10 U or 100 U of rhIL-2. \*p < 0.05.

Among the natural ligands of PPAR $\alpha$ , various eicosanoids, such as leucotriene B<sub>4</sub>, are generated during inflammatory responses [20, 21]. In addition to these effects of ligand-bound PPAR $\alpha$  on NF- $\kappa$ B and c-jun, it has also been reported that ligand-free PPAR $\alpha$  may inhibit the phosphorylation of p38 [22], a kinase that also promotes IFN $\gamma$  transcription downstream of TCR stimulation [23].

Thus, we analyzed the *in vitro* response of human V $\gamma$ 9V $\delta$ 2 T cells stimulated with a PPAR $\alpha$  agonist, LY171883. Early activation was measured by the surface expression of CD69 by V $\gamma$ 9V $\delta$ 2 T cells after 2 days of fresh PBMC culture with BrHPP. Among the living lymphocytes, gated TCR V $\gamma$ 9<sup>+</sup> cells expressed CD69 if activated by BrHPP (Fig. 2A). Preincubation with LY-171883 led to a significant dose-dependent decrease in this expression (Fig. 2A and B).

Then, we analyzed the effect of PPAR $\alpha$  agonists on the intracellular expression of IFN $\gamma$  by V $\gamma$ 9V $\delta$ 2 T cells. BrHPP-activation with a blocking secretion led to an expression of intracellular IFN $\gamma$ , which was inhibited by the preincubation with LY-171883 in a dose-dependent manner (Fig. 2C and D). Finally, we showed that LY-171883 induced an inhibition of the V $\gamma$ 9V $\delta$ 2 T cell proliferation, evaluated by the CFSE dilution, when cultured for six days with a weak or high dose of rhIL-2 (Fig. 2E). Thus, *in vitro* activation of the PPAR $\alpha$  pathway leads to the inhibition of BrHPP-induced activation and proliferation of human V $\gamma$ 9V $\delta$ 2 T cells.

We know from *in vivo* studies that V $\gamma$ 9V $\delta$ 2 T cells from macaques respond to a first BrHPP injection by amplification in the blood, but they become non-responsive following several BrHPP injections [7, 8]. Our finding, which highlights the activation of the PPAR $\alpha$  pathway in V $\gamma$ 9V $\delta$ 2 T cells from primates injected three times with BrHPP, can be thus associated with the non-response of these cells under the same *in vivo* experimental conditions. In humans, repeated exposure to BrHPP also leads to rapid exhaustion of the V $\gamma$ 9V $\delta$ 2 T cell response [9]. Moreover, the similarity of human and non-human primate V $\gamma$ 9V $\delta$ 2 T cells allows the correlation of the two discoveries. With this study, we highlighted the involvement of the PPAR $\alpha$  pathway in the antigen-selective desensitization of human V $\gamma$ 9V $\delta$ 2 T cells.

The use of inhibitors of the PPAR $\alpha$  pathway or a method preventing its activation could enhance the advantage of using cancer immunotherapy based on V $\gamma$ 9V $\delta$ 2 T cells. Indeed, the repression of PPAR $\alpha$  expression can be induced by TNF $\alpha$  [24] or by a group of N-(methylsulfonyl)amides derived from PPAR $\alpha$  agonist carboxylic acids with an antagonist behavior on PPAR $\alpha$  [25]. Some kinases or their activators can also lead to specific inhibition of PPAR $\alpha$  transcriptional activity, such as the MEK1 or AMP-activated protein kinase activators 5-aminoimidazole-4-carboxamide riboside [26, 27]. Furthermore, it was shown very recently that PPAR $\alpha$  inhibition could improve the therapeutic efficacy of glucocorticoids in aggressive forms of chronic lymphocytic leukemia [28]. Thus, PPAR $\alpha$  inhibition could be used in V $\gamma$ 9V $\delta$ 2 T cell-based cancer immunotherapy.

**Acknowledgements.** We would like to thank Laurence Lamant and Pierre Brousset for their helpful suggestions. Innate Pharma graciously provided us with clinical grade batches of BrHPP. Thank you also to Laboratoires Sanofi-Aventis for the kind gift of recombinant human IL-2. This study was supported in part by institutional grants from the Institut National de la Santé et de la Recherche Médicale (INSERM) and the Université de Toulouse 3 and contracts from Institut National du Cancer (RITUXOP and V9V2TER).

**Conflict of interest.** The authors declare no relevant conflict of interest.

## REFERENCES

1. Angelini, D.F., Borsellino, G., Poupot, M., Diamantini, A., Poupot, R., Bernardi, G., Poccia, F., Fournie, J. J. and Battistini, L. FcγRIII discriminates between 2 subsets of Vγ9Vδ2 effector cells with different responses and activation pathways. **Blood** 104 (2004) 1801-1807.
2. Rothenfusser, S., Buchwald, A., Kock, S., Ferrone, S. and Fisch, P. Missing HLA class I expression on Daudi cells unveils cytotoxic and proliferative responses of human γδ T lymphocytes. **Cell. Immunol.** 215 (2002) 32-44.
3. Kunzmann, V. and Wilhelm, M. Anti-lymphoma effect of γδ T cells. **Leuk. Lymphoma** 46 (2005) 671-680.
4. Bonneville, M. and Scotet, E. Human Vγ9Vδ2 T cells: promising new leads for immunotherapy of infections and tumors. **Curr. Opin. Immunol.** 18 (2006) 539-546.
5. Kabelitz, D., Wesch, D. and He, W. Perspectives of γδ T cells in tumor immunology. **Cancer Res.** 67 (2007) 5-8.
6. Caccamo, N., Meraviglia, S., Cicero, G., Gulotta, G., Moschella, F., Cordova, A., Gulotta, E., Salerno, A. and Dieli, F. Aminobisphosphonates as new weapons for γδ T cell-based immunotherapy of cancer. **Curr. Med. Chem.** 15 (2008) 1147-1153.
7. Sicard, H., Ingoure, S., Luciani, B., Serraz, C., Fournie, J.J., Bonneville, M., Tiollier, J. and Romagne, F. In vivo immunomanipulation of Vγ9Vδ2 T cells with a synthetic phosphoantigen in a preclinical nonhuman primate model. **J. Immunol.** 175 (2005) 5471-5480.
8. Cendron, D., Ingoure, S., Martino, A., Casetti, R., Horand, F., Romagne, F., Sicard, H., Fournie, J.J. and Poccia, F. A tuberculosis vaccine based on phosphoantigens and fusion proteins induces distinct γδ and αβ T cell responses in primates. **Eur. J. Immunol.** 37 (2007) 549-565.
9. Sicard, H., Rossi, J.P., Rousselot, P., Colette, A., Paiva C., Ingoure, S. and Lafaye de Micheaux, S. Cognitive study of reactivity to IPH1101 of peripheral γδ T lymphocytes from chronic myeloid leukemia, multiple myeloma and follicular lymphoma patients. **ASH Annual Meeting Abstracts** 112 (2008) 1530.

10. Barbier, O., Torra, I.P., Duguay, Y., Blanquart, C., Fruchart, J.C., Glineur, C. and Staels, B. Pleiotropic actions of peroxisome proliferator-activated receptors in lipid metabolism and atherosclerosis. **Arterioscler. Thromb. Vasc. Biol.** 22 (2002) 717-726.
11. Daynes, R.A. and Jones, D.C. Emerging roles of PPARs in inflammation and immunity. **Nat. Rev. Immunol.** 2 (2002) 748-759.
12. Delerive, P., De Bosscher, K., Besnard, S., Vanden Berghe, W., Peters, J.M., Gonzalez, F. J., Fruchart, J.C., Tedgui, A., Haegeman, G. and Staels, B. Peroxisome proliferator-activated receptor alpha negatively regulates the vascular inflammatory gene response by negative cross-talk with transcription factors NF-kappaB and AP-1. **J. Biol. Chem.** 274 (1999) 32048-32054.
13. Zhang, M.A., Rego, D., Moshkova, M., Kebir, H., Chruscinski, A., Nguyen, H., Akkermann, R., Stanczyk, F.Z., Prat, A., Steinman, L. and Dunn, S.E. Peroxisome proliferator-activated receptor (PPAR)alpha and -gamma regulate IFNgamma and IL-17A production by human T cells in a sex-specific way. **Proc. Natl. Acad. Sci. USA** 109 (2012) 9505-9510.
14. Martinet, L., Poupot, R. and Fournie, J.J. Pitfalls on the roadmap to gammadelta T cell-based cancer immunotherapies. **Immunol. Lett.** 124 (2009) 1-8.
15. Capietto, A.H., Martinet, L. and Fournie, J.J. How tumors might withstand gammadelta T-cell attack. **Cell. Mol. Life Sci.** 68 (2011) 2433-2442.
16. Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S. and Mesirov, J.P. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. **Proc. Natl. Acad. Sci. USA** 102 (2005) 15545-15550.
17. Boullier, S., Poquet, Y., Debord, T., Fournie, J.J. and Gougeon, M.L. Regulation by cytokines (IL-12, IL-15, IL-4 and IL-10) of the Vgamma9Vdelta2 T cell response to mycobacterial phosphoantigens in responder and anergic HIV-infected persons. **Eur. J. Immunol.** 29 (1999) 90-99.
18. Martini, F., Paglia, M.G., Montesano, C., Enders, P.J., Gentile, M., Pauza, C.D., Gioia, C., Colizzi, V., Narciso, P., Pucillo, L.P. and Poccia, F. V gamma 9V delta 2 T-cell anergy and complementarity-determining region 3-specific depletion during paroxysm of nonendemic malaria infection. **Infect. Immun.** 71 (2003) 2945-2949.
19. Montesano, C., Gioia, C., Martini, F., Agrati, C., Cairo, C., Pucillo, L.P., Colizzi, V. and Poccia, F. Antiviral activity and anergy of gammadeltaT lymphocytes in cord blood and immuno-compromised host. **J. Biol. Regul. Homeost. Agents** 15 (2001) 257-264.
20. Luna-Gomes, T., Bozza, P.T. and Bandeira-Melo, C. Eosinophil recruitment and activation: the role of lipid mediators. **Front. Pharmacol.** 4 (2013) 27.

21. Montuschi, P. LC/MS/MS analysis of leukotriene B4 and other eicosanoids in exhaled breath condensate for assessing lung inflammation. **J. Chromatogr. B. Analyt. Technol. Biomed. Life Sci.** 877 (2009) 1272-1280.
22. Hou, X., Shen, Y.H., Li, C., Wang, F., Zhang, C., Bu, P. and Zhang, Y. PPARalpha agonist fenofibrate protects the kidney from hypertensive injury in spontaneously hypertensive rats via inhibition of oxidative stress and MAPK activity. **Biochem. Biophys. Res. Commun.** 394 (2010) 653-659.
23. Mendler, A.N., Hu, B., Prinz, P.U., Kreutz, M., Gottfried, E. and Noessner, E. Tumor lactic acidosis suppresses CTL function by inhibition of p38 and JNK/c-Jun activation. **Int. J. Cancer** 131 (2012) 633-640.
24. Lim, W.S., Ng, D.L., Kor, S.B., Wong, H.K., Tengku-Muhammad, T.S., Choo, Q.C. and Chew, C.H. Tumour necrosis factor alpha down-regulates the expression of peroxisome proliferator activated receptor alpha (PPARalpha) in human hepatocarcinoma HepG2 cells by activation of NF-kappaB pathway. **Cytokine** 61 (2013) 266-274.
25. Ammazalorso, A., D'Angelo, A., Giancristofaro, A., De Filippis, B., Di Matteo, M., Fantacuzzi, M., Giampietro, L., Linciano, P., Maccallini, C. and Amoroso, R. Fibrate-derived N-(methylsulfonyl)amides with antagonistic properties on PPARalpha. **Eur. J. Med. Chem.** 58 (2012) 317-322.
26. el Azzouzi, H., Leptidis, S., Bourajjaj, M., van Bilsen, M., da Costa Martins, P.A. and De Windt, L.J. MEK1 inhibits cardiac PPARalpha activity by direct interaction and prevents its nuclear localization. **PLoS One** 7 (2012) e36799.
27. Sozio, M.S., Lu, C., Zeng, Y., Liangpunsakul, S. and Crabb, D.W. Activated AMPK inhibits PPAR- $\alpha$  and PPAR- $\gamma$  transcriptional activity in hepatoma cells. **Am. J. Physiol. Gastrointest. Liver Physiol.** 301 (2011) G739-747.
28. Tung, S., Shi, Y., Wong, K., Zhu, F., Gorczynski, R., Laister, R.C., Minden, M., Blechert, A.K., Genzel, Y., Reichl, U. and Spaner, D.E. PPARalpha and fatty acid oxidation mediate glucocorticoid resistance in chronic lymphocytic leukemia. **Blood** 122 (2013) 969-980.